

# Comet™



## Economical Post-Emergent Control of Broadleaf Weeds.

Comet™ is a selective post-emergence herbicide that controls annual and perennial broadleaf weeds and woody brush in rangeland and permanent pastures. Comet may also be used for selective post-emergence control of annual and perennial broadleaf weeds and volunteer potatoes in small grains, field corn, sweet corn, grain sorghum, fallow cropland, on-farm non-cropland and pome fruit.

Comet may be applied as a single broadcast treatment or spot treatment to control susceptible broadleaf weeds in on-farm non-cropland areas such as fence rows, building perimeters, around irrigation equipment and on-farm private roadways.

### KEY PERFORMANCE BENEFITS

- Excellent control of broadleaf weeds
- Can be tank-mixed with other broadleaf and grass herbicides as well as most fungicides, insecticides and foliar fertilizers
- Offers rotational flexibility to next year's crop

EPA REG. NO. 71368-87

ACTIVE INGR. fluroxypyr (26.2%)

FORMULATION liquid

CHEM. FAMILY carboxylic acid

HRAC NO. 4

MODE OF ACT. synthetic auxin growth regulator

REI 12 hours

SIGNAL WORD warning

RESTRICTED no

PACKAGE SIZE 2 x 2.5 gal, 30 gal, 265 gal tote



Nufarm

Grow a better tomorrow.

# Comet™

## KEY USES

Rangeland  
 Permanent pastures  
 Small grains, field corn, sweet corn,  
 grain sorghum, fallow cropland,  
 and on-farm non-cropland  
 Pome fruit

## KEY WEEDS CONTROLLED

*(See product label for complete list)*

Bedstraw (*cleavers*)  
 Bindweed, field\*  
 Bindweed, hedge  
 Buckwheat, wild\*  
 Canola, volunteer\*  
 Chickweed  
 Clover, white  
 Cocklebur  
 Coffeeweed  
 Devilsclaw\*  
 Flax, volunteer  
 Grape spp.  
 Hemp dogbane  
 Horsetail, field\*  
 Horseweed (*maretail*)\*  
 Jimsonweed  
 Knotweed\*  
 Kochia\*  
 Mallow, common\*  
 Mallow, Venice  
 Marshelder\*  
 Morningglory  
 Mustard spp.\*  
 Nightshade spp.\*  
 Pennycress, field\*  
 Potato, volunteer\*  
 Prickly lettuce  
 Puncturevine  
 Purslane, common  
 Ragweed, common  
 Sunflower  
 Thistle, Russian\*  
 Velvetleaf

\* Suppression

## USE SUMMARY

Avoid unintentional spray drift to adjacent susceptible broadleaf plants or broadleaf crops.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Use application rates of 2/3 to 1 1/3 pints per acre depending on crop (2/3 to 2 2/3 for pome fruit). One application is allowed per season.

*See label for complete directions for use.*

