

Maestro[®] 2EC



Flexible, Fast-Acting Broadleaf Control.

Maestro[®] 2EC is a 33.4% bromoxynil formulation that provides control of a variety of difficult broadleaf weeds in corn, sorghum, wheat, non-crop and many other crops and uses.

EXCELLENT COMPONENT OF HERBICIDE RESISTANCE MANAGEMENT PROGRAM

- Controls more than 40 tough broadleaf weeds
- Compatible with many grass and broadleaf herbicides
- Flexible application options
- Over 45 years of use with few resistance issues

EPA REG. NO.	71368-29
ACTIVE INGR.	bromoxynil octanoate (33.4%)
FORMULATION	liquid
CHEM. FAMILY	nitrile
HRAC NO.	6
MODE OF ACT.	inhibition of photosynthesis
REI	24 hours
SIGNAL WORD	warning
RESTRICTED	no
PACKAGE SIZE	2 x 2.5 gal



Grow a better tomorrow.

Maestro® 2EC

KEY USES

Alfalfa (*seedling*)
 Barley
 Corn (*field and popcorn*)
 Flax
 Garlic
 Mint
 Oats
 Onion (*dry bulb*)
 Rye
 Sorghum (*grain and forage*)
 Triticale

KEY WEEDS CONTROLLED

(See product label for complete list)

Black nightshade
 Bristly starbur
 Buckwheat, wild
 Buffalobur
 Cocklebur, common
 Cutleaf nightshade
 Eastern black nightshade
 Groundsel, common
 Hairy nightshade
 Hemp sesbania
 Jimsonweed
 Knawel
 Kochia
 Lambsquarters, common
 London rocket
 Mayweed
 Morningglory; ivyleaf, tall
 Mustard, blue
 Pennycress, field
 Pepperweed (*annual*)
 Pigweed; redroot, spiny
 Puncturevine
 Ragweed; common, giant
 Shepherd's-purse
 Smartweed; giant, Pennsylvania
 Sowthistle, annual
 Sunflower
 Thistle, Russian
 Velvetleaf

USE SUMMARY

This product is formulated as an emulsifiable concentrate of octanoic acid ester of bromoxynil containing the equivalent of 2 pounds of bromoxynil per gallon.

Optimum weed control is obtained when this product is applied to actively growing weed seedlings. This product is primarily a contact herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control.

Maestro® 2EC Alone: Fill the tank 1/2 to 3/4 full with clean water. Begin agitation and add product. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

Tank Mixtures: This product can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. To apply this product in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with a wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add this product and water to the spray tank to the desired level. If tank mixing with other product types, add this product first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

ROTATION RESTRICTIONS

Do not plant rotational crops within 30 days following application of Maestro 2EC.

See label for complete directions for use..

