Maestro® 4EC



Flexible, Fast-Acting Broadleaf Control.

Maestro® 4EC offers growers excellent control of more than 50 tough broadleaf weeds including cocklebur, common lambsquarters, morningglory and velvetleaf. This fast-acting post-emergence herbicide can be used as a stand alone product or in tank-mix combinations with broadleaf herbicides, grass herbicides, insecticides, plant growth regulators or spray grade liquid fertilizers.

PROPRIETARY FORMULA FOR EXCELLENT CONTROL

- Controls more than 50 tough broadleaf weeds
- Concentrated mixed-ester formulation of bromoxynil
- Flexible application window
- Can be applied by ground or aerial application or through sprinkler irrigation equipment
- Compatible with numerous other grass and broadleaf active ingredients
- Proven active ingredients in quality formulation
- Rotational crops may be planted 30 days after application

EPA REG. NO.	71368-78
ACTIVE INGR.	bromoxynil (55%)
FORMULATION	liquid
CHEM. FAMILY	nitrile
GROUP NO.	6
MODE OF ACT.	inhibition of photosynthesis
MODE OF ACT. REI	inhibition of photosynthesis 24 hours
	. ,
REI	24 hours
REI SIGNAL WORD	24 hours warning

Maestro® 4EC

KEY USES

Alfalfa (seedling)

Barley

Corn (field and popcorn)

Flax Garlic Mint

Oats

Onion (dry bulb)

Rye

Sorghum (grain and forage)

Triticale

KEY WEEDS CONTROLLED

(See product label for complete list)

Annual sowthistle

Black nightshade

Blue mustard

Bristly starbur

Buffalobur

Cocklebur, common

Cutleaf nightshade

Devil's claw

Eastern black nightshade

Field pennycress

Groundsel, common

Hairy nightshade

Hemp sesbania

Hophornbean copperleaf

Jimsonweed

Knawel

Kochia

Lambsquarters, common

London rocket

Mayweed

Morningglory; ivyleaf, pitted

Mustard, tumble

Pepperweed (annual)

Pigweed; redroot, spiny

Prairie sunflower

Puncturevine

Ragweed; common, giant

Shepherd's-purse

Smartweed; green, Pennsylvania

Sunflower Thistle, Russian Velvetleaf

USE SUMMARY

Post-emergence application of Maestro® 4EC herbicide will control weeds when sprayed in the seedling stage up to 8" in height depending on species, timing and application rate.

RATES

Stand-alone: 1/2 to 1 pint/A Tank-mix: 3/8 to 1 pint/A

ROTATION RESTRICTIONS

Do not plant rotational crops within 30 days following application of Maestro 4EC.

OTHER INFORMATION

Maestro 4EC is primarily a contact herbicide; therefore thorough coverage of the weed seedlings is essential for optimum control.

Maestro 4EC has little residual activity, and subsequent flushes of weeds will not be controlled by the initial treatment.

See label for complete directions for use.

