

Maestro™ D

Selective Herbicide



HERBICIDE



Post-Emergent Selective Herbicide

Active Ingredient:

2.0 lbs/gal Bromoxynil

1.9 lbs/gal 2,4-D

FEATURES & BENEFITS

- Convenient premix formulation including a contact and systemic herbicide.
- Provides control of a variety of difficult broadleaf weeds including wild buckwheat, redroot pigweed and wild mustard.
- Ground and aerial application options.
- Quick weed knock down.

PRIMARY WEEDS CONTROLLED

Broadleaf Weeds in Wheat, Barley, Oats, Triticale and Rye.

PROFILE

Maestro D is a premix of bromoxynil and 2,4-D ester that provides selective post-emergent control of important broadleaf weeds infesting the small grains crops wheat, barley, oats, rye and triticale.

ADVANTAGE

Optimum weed control is obtained when Maestro D is applied to actively growing weed seedlings. Maestro D contains a contact herbicide and a systemic herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control.

MODE OF ACTION

Maestro D is formulated as an emulsifiable concentrate of octanoic acid ester of bromoxynil containing the equivalent of 2 pounds of bromoxynil per gallon and 2,4- Dichlorophenoxyacetic acid containing the equivalent of 1.9 pounds per gallon.

Maestro is a trademark of Nufarm Americas, Inc.

Always follow directions found on container label. This information is for promotional purposes only.



Nufarm Agriculture Division

www.ag.us.nufarm.com

GENERAL INFORMATION

Application Rates

Use 3/4 - 1 pint of Maestro D per acre.

Ground Application - In general, a spray volume of 10 to 20 gallons per acre is recommended for optimum spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles with a spray pressure of 40-60 psi are recommended.

Aerial Application - In general, a minimum spray volume of 5 gallons per acre and a maximum pressure of 40 psi are recommended.

*** Application should be made from the fully tillered stage of growth but before the jointing stage.

*** The 1 pint rate should be used for mustards, pigweed, wild buckwheat, kochia or when weeds are growing under dry cool conditions.

| WEEDS CONTROLLED | | | |
|---|----------------|----------------|------------------|
| Apply to weeds up to the 4 leaf stage or 2 inches in height | | | |
| Mustards | Field Bindweed | Wild Buckwheat | Cow Cockle |
| Frenchweed | Gumweed | Plantain | Russian Thistle |
| Nightshade | Pigweed | Sunflower | Annual Smartweed |
| Arrowhead | Ragweed | Canola | Lambsquarter |

Useful Information:

- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 45 days after treatment
- Do not plant rotational crops within 30 days following application
- Total cumulative rate should not exceed 2 lbs. a.i. of 2,4-D
- Do not feed treated straw to livestock
- Personal Protective Equipment (PPE) - Applicators and handlers must wear long-sleeved shirt and pants, chemical resistant gloves, protective eyewear, shoes and socks, and a chemical resistant apron when cleaning equipment.
- Spray drift management practices such as controlling droplet size, considering boom length and height, and monitoring wind, temperature and humidity, are important to prevent off-target plant injuries



Nufarm Americas Inc.
150 Harvester Drive • Suite 200
Burr Ridge, IL 60527

Phone: 630.455.2000 • Fax: 630.455.2001
www.ag.us.nufarm.com

PIB.0107.Maestro D
© 2007 Nufarm Americas Inc.

Always follow directions found on container label.
This information is for promotional purposes only.