

26/36®



Dual Mode of Action for Effective Disease Control.

Incorporating 26/36® fungicide into an integrated disease management program provides effective control of key diseases of turfgrass and ornamentals.

EASY-TO-USE FORMULA

- Easy-to-use, premixed formulation
- Convenient
- Broad-spectrum disease control for turf
- Tank-mix compatible with most commonly used pesticides

EPA REG. NO. 228-630

ACTIVE INGR. iprodione (19.65%),
thiophanate-methyl (19.65%)

FORMULATION flowable

CHEM. FAMILY dicarboxamide, benzimidazole

FRAC NO. 1 and 2

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 2 x 2.5 gal

 COMMERCIAL

 TURF

 LANDSCAPE ORNAMENTALS

 GOLF TURF

 SOD FARMS



Nufarm

Grow a better tomorrow.

26/36®

KEY USES

Golf courses
 Institutional areas
 Ornamentals
 Sod farms
 Turfgrass

KEY DISEASES CONTROLLED

(See product label for complete list)

Anthracnose
 Ascochyta leaf blight
 Brown patch
 Copper spot
 Corticium
 Dollar spot
 Fusarium patch
 Gray snow mold
 Leaf spot
 Pink snow mold
 Red thread
 Zoysia patch

USE SUMMARY

26/36® fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. It provides effective, broad spectrum turf disease control and also serves as a useful tank mixture in the resistance management program required for other resistant sensitive fungicides.

When used in conjunction with good turf management practices, 26/36 Fungicide is effective in controlling the following diseases: dollar spot, brown patch, anthracnose, leaf spots such as Helminthosporium leaf spot caused by Drechslera spp. pathogens, corticium red thread, Fusarium patch, zoysia patch, ascochyta leaf blight and copper spot.

Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

See label for complete directions for use.

APPLICATION PRECAUTIONS AND RESTRICTIONS

DO NOT exceed a total of 14.5 fl oz product per 1000 sq ft per year.

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. When disease pressure is low to moderate, the lower rates and longer intervals are recommended.

DO NOT mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred.

DO NOT mix with any sticker, extender, or wetting agent.

DO NOT graze animals on treated turf.

DO NOT feed clippings from treated turf to livestock or poultry.

See label for complete application precautions and restrictions.

