

This seed has been treated with metalaxyl and fludioxonil fungicide (Spirato M 185 FS) seed treatment at rates specified by the manufacturer.

CAUTION

TREATED SEED

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES. STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS.

KEEP OUT OF REACH OF CHILDREN

DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Treated seed may be harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Seed handlers: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Seed Handlers must wear:

- Long pants, long-sleeved shirt, and protective gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Treated seed exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting. **DO NOT** contaminate water bodies when disposing of planting equipment wash water.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

RESTRICTIONS:

After the seeds have been planted, do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 24 hours**. Exception: If the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

Forage may not be grazed until 30 days after planting.

PACKAGE AND SEED DISPOSAL:

Dispose of all excess treated seed by burying seed away from bodies of water. **DO NOT** contaminate bodies of water when disposing of planting equipment wash water. Dispose of seed packaging or containers in accordance with local requirements. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 042616

13841000

Spirato™ is a trademark of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

