

# Nuprid® 2SC

INSECTICIDE

## PROVEN, BROAD-SPECTRUM INSECT CONTROL.

Nuprid® 2SC systemic insecticide provides economical and long-lasting control of a broad spectrum of damaging insects. The expanded label allows for both soil and foliar application on many different crops. It offers outstanding control of aphids, whiteflies, mealybugs, Colorado potato beetles, leafhoppers and many other sucking and piercing pests. Plus, Nuprid 2SC can help stop transmission of diseases caused by insect feeding. Nuprid 2SC delivers consistent field performance, plus it's easy to mix, use and apply.

### COMPETITIVE ADVANTAGES

- Selective control of key sucking and piercing insects
- Expanded label allows for soil and foliar application on multiple crops
- Reliable performance against yield robbing pests
- Helps stop transmission of many insect-vectored diseases
- Readily mixes with other tank-mix partners to save a trip across the field

### AT-A-GLANCE INFO

EPA REG. NO.	228-572
ACTIVE INGR.	imidacloprid (21.4%)
FORMULATION	liquid
CHEM. FAMILY	neonicotinoid
GROUP NO.	4A
MODE OF ACT.	nicotinic acetylcholine receptor agonists
REI	12 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	1 gal (4 per case), 265 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Grapple  | Phostrol® | Tuscany SC 

 **Nufarm**  
Grow a better tomorrow

# Nuprid® 2SC

PROVEN, BROAD-SPECTRUM INSECT CONTROL.

## KEY USES

Citrus  
Cotton  
Fruiting vegetables  
Grapes  
Leafy green vegetables  
Legume vegetables  
Pome fruits  
Potatoes  
Soybeans  
Stone fruits  
Strawberries  
Tree, bush and vine crops  
Tree nuts  
Tropical fruits

## KEY INSECTS CONTROLLED

(See product label for complete list)

Aphids  
Black fly  
Cucumber beetles  
Flea beetles  
Japanese beetle  
Leafhoppers  
Leafminers  
Mealybugs  
Mole crickets  
Plant bugs  
Psyllids  
Sharpshooters  
Thrips  
Whiteflies

## KEY DISEASES CONTROLLED

(See product label for complete list)

Bacterial wilt  
Bean common mosaic virus  
Beet curly top hybrigeminivirus  
Citrus yellows  
Net necrosis  
Pecan scab  
Potato yellows  
Tomato mottle virus

## USE SUMMARY

### APPLICATION RATES

**Foliar Application:** Apply as a directed or broadcast foliar spray. Thorough coverage of foliage is necessary for optimum control. Apply with properly calibrated ground or aerial application equipment at a minimum of 10 gals/A by ground or 5 gals/A by air unless otherwise specified on crop-specific application sections.

**Soil Application:** Make applications into the seed or root-zone of crop. Nuprid® 2SC may be applied with ground or chemigation equipment. Best activity results from applications to the root-zone of plants to be protected.

The earlier Nuprid 2SC is available to developing plants, the earlier protection begins. Nuprid 2SC is continuously taken into the roots over a long period of time and its systemic nature allows movement from roots through the xylem tissue to all vegetative parts of the plant, resulting in extended residual activity, insect control and prevention and reduction of virus transmission. Higher rates are recommended when infestations occur later in crop development or where pest pressure is continuous.

Do not apply more than 0.5 lbs active ingredient per acre per crop season, regardless of method of application, unless specified within a crop-specific application section.

See product label for complete directions for use.



[nufarm.com/uscrop](http://nufarm.com/uscrop)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/uscrop](http://www.nufarm.com/uscrop) to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. Grapple®, Nuprid®, Phostrol® and Tuscany® are trademarks of Nufarm.



15-AG-0031-C