



Grapple™

SELECTIVE HERBICIDE

EXCELLENT WEED CONTROL WITH RESIDUAL.

Grapple™ is a selective herbicide for burndown and residual control of many annual grass and broadleaf weeds in a variety of crop timings including pre- and post-emerge field corn and tomatoes, pre-plant soybeans and cotton, and fallow land. Count on Grapple to keep crops weed-free with lasting residual. Plus, Grapple is tank mix compatible with many pre-emerge herbicides for enhanced weed control.

COMPETITIVE ADVANTAGES

FORMULATED FOR WEED CONTROL IN MANY CROPS

- Outstanding broad-spectrum pre- and post-emergent control of many weeds
- Labeled for use on a wide range of crops
- Long-lasting residual control
- Tank mix compatible with many other herbicides for added weed control

AT-A-GLANCE INFO

EPA REG. NO.	71368-121
ACTIVE INGR.	rimsulfuron (25%)
FORMULATION	water-dispersible granule
CHEM. FAMILY	sulfonylurea
GROUP NO.	2
MODE OF ACT.	ALS inhibitor (acetolactate synthase)
REI	4 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	20 oz bag (12 per case)

MORE PRODUCTS EXCLUSIVELY FROM NUFARM



Nufarm
Grow a better tomorrow



EXCELLENT WEED CONTROL WITH RESIDUAL.

KEY USES

Field corn
Field-grown tomatoes
Pre-plant soybeans and cotton
Also labeled for use on blueberry (high and low bush)*, caneberry (raspberry, blackberry)*, citrus fruit, grapes, potatoes, potatoes grown for seed, stone and pome fruit, tree nuts

KEY WEEDS CONTROLLED

(See product label for complete list)

PRE-EMERGE

Barnyardgrass
Chamomile, false
Crabgrass, large (partial)
Dandelion, common
Filaree, redstem
Foxtail; giant, green, yellow
Henbit
Kochia
Mustard; birdsrape, black
Pigweed; redroot, smooth
Purslane, common

POST-EMERGE

Grassy: 1" to 2" / Broadleaf: 1" to 3"

Barnyardgrass
Bluegrass, annual
Chamomile, false
Chickweed, common
Crabgrass, large (½")
Foxtail; bristly, giant, green, yellow
Henbit
Kochia
Mustard; birdsrape, black, wild
Panicum, fall
Pigweed; redroot, smooth
Puncturevine
Purslane, common
Shepherd's-purse
Wild radish

*Not for use in California

USE SUMMARY

PRE- AND POST-EMERGE TO FIELD CORN (EXCEPT CALIFORNIA*)

Fallow (burndown) applications: Apply Grapple at 1 to 2 oz/A. Grapple may be used as a fallow treatment in the spring or fall when the majority of weeds have emerged and are actively growing. Field corn may be planted to this treated area at any time.

Pre-emerge applications: Apply Grapple at 0.5 to 2 oz/A before corn emergence. Nufarm specifies a rate of 1 to 1.5 oz/A for most applications. Grapple may be applied pre-emergence or preplant to corn. Applications of Grapple made before weed emergence will provide residual control of labeled weeds. Control of emerged weeds will require the addition of spray adjuvants as noted on the label.

Post-emerge applications: Grapple may be applied at 0.5 to 2 oz/A as a post-emerge broadcast application. Nufarm specifies a rate of 1 oz/A for most applications. Apply Grapple to corn that is up to 12 inches tall. Do not apply to corn taller than 12 inches or exhibiting 6 or more leaf collars, whichever is more restrictive. Applications of Grapple made after weed emergence will provide contact control of labeled weeds as well as limited residual control of later emergence. For control of emerged weeds, application of Grapple must include a nonionic surfactant and an ammonium nitrogen fertilizer.

PREPLANT TO COTTON AND SOYBEAN: Grapple may be applied preplant after fall harvest through early spring 30 days or more prior to planting, whenever the ground is not frozen, to control emerged weeds and to provide limited residual control of early-emerging spring weeds. Apply Grapple at 1 oz/A.

Grapple may be tank mixed with other herbicides registered for preplant cotton and soybean including glyphosate, paraquat, glufosinate, 2,4-D, and dicamba. Always follow directions of the most restrictive label.

DIRECT-SEEDED OR TRANSPLANT TOMATOES (EXCEPT CALIFORNIA*)

Pre-emerge applications: Apply Grapple at 2 to 4 oz/A. To activate Grapple in the soil, supply moisture by a single rainfall event, or apply sprinkler irrigation of 1/2 to 1 inch within 5 days after application.

Post-emerge applications: Apply Grapple at 1 to 2 oz/A to young, actively growing weeds after the crop has reached the cotyledon stage. Optimum performance is obtained when weeds are less than 1 inch in height or diameter and are actively growing. Use a surfactant at a minimum rate of 2.25% V/V. For best results, 1/2 to 1 inch of rainfall or sprinkler irrigation should be applied no sooner than 4 hours and no more than 5 days after application of Grapple.

See label for complete directions for use.

*See label for complete directions for use when used in California field corn and tomato crops.

PRECAUTIONS AND RESTRICTIONS

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain, or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water that may be used on other crops.
- Carefully observe sprayer cleanup instructions, as spray tank residue may damage crops other than potatoes or tomatoes.

See label for complete application precautions and restrictions.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. Cheelah®, Duplosan™, Grapple™, and Panther® are trademarks of Nufarm.



19-AG-0009