

# Spyder® Extra

SELECTIVE HERBICIDE

## DUAL-ACTION CONTROL FOR CONIFER PLANTATIONS & NON-CROP AREAS.

Spyder® Extra contains two active ingredients – sulfometuron and metsulfuron – for outstanding foliar and residual control of dozens of annual and perennial grasses and broadleaf weeds, as well as sedges. Spyder Extra offers multiple use rates so you can tailor applications to specific needs, and its high-quality, low-rate formulation makes it easy to mix and apply.

### COMPETITIVE ADVANTAGES

- Provides broad-spectrum control of annual and perennial grasses and broadleaf weeds
- Offers foliar and residual control for selective applications to non-crop sites
- Low use rates
- Easy-to-use granular formulation
- Excellent tank-mix partner for Razor® Pro, WEEDestroy® AM-40 and Veteran® 720

### AT-A-GLANCE INFO

EPA REG. NO.	228-690
ACTIVE INGR.	sulfometuron-methyl (56.25%), metsulfuron methyl (15%)
FORMULATION	water-dispersible granule
CHEM. FAMILY	sulfonylurea
GROUP NO.	2
MODE OF ACT.	ALS inhibitor
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	4 x 5 lb

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Clipper® SC | Veteran® 720 | Lock Down® SC

 **Nufarm**  
Grow a better tomorrow

# Spyder® Extra

## DUAL-ACTION CONTROL FOR CONIFER PLANTATIONS & NON-CROP AREAS.

### KEY USES

Conifer plantations  
Non-crop sites  
Unimproved turf grasses

### KEY WEEDS CONTROLLED

(See product label for complete list)

Aster	Lambsquarters
Bahiagrass	Miners lettuce
Barnyardgrass	Mouseear chickweed
Black mustard	Multiflora rose (wild roses)
Blackberry	Musk thistle
Black-eyed Susan	Oxeye daisy
Blue mustard	Pennsylvania smartweed
Chickweed	Pepperweed
Clover	Pigweed
Cocklebur	Plantain
Crabgrass	Ragweed
Curly dock	Salsify
Dandelion	Shepherd's-purse
Dogfennel	Signalgrass
Downy brome (cheat)	St. Johnswort
Dyer's woad	Tansy mustard
Evening primrose	Teasel
Fescue	Vetch
Fireweed	Whitetop
Fleabanes	Woodsorrel
Foxtail	Yankee weed
Goldenrod	Yellow nutsedge
Henbit	Yellow rocket
Honeysuckle	
Johnsongrass	

### USE SUMMARY

#### FORESTRY

##### USE RATES BEFORE TRANSPLANTING

Conifer site preparation: use rates for selected species before transplanting conifers.

Loblolly Pine	3 to 4 oz/A When to plant into treated areas: Planting season following application.
Slash Pine	3 to 4 oz/A When to plant into treated areas: Planting season following application.
Black Spruce	2 2/3 to 5 1/3 oz/A When to plant into treated areas: Not less than 13 months following application.
Red Pine	1 1/3 to 2 2/3 oz/A When to plant into treated areas: The following spring or summer but not less than 3 months after application. Areas receiving 2/3 to 1 1/3 oz/A may be transplanted in a minimum of 30 days following application.
Douglas Fir	2 2/3 to 5 1/3 oz/A When to plant into treated areas: Planting season following application.

##### USE RATES AFTER TRANSPLANTING

Conifer release: use rates for selected species after transplanting conifers to control certain species of hardwoods, broadleaf weeds and grasses.

Loblolly Pine	2 2/3 to 4 oz/A
Slash Pine	2 2/3 to 3 oz/A

##### NON-AGRICULTURAL USES

Non-crop Areas	2 2/3 to 5 1/3 oz/A depending on target weeds. Use the higher rate for areas with heavy weed growth, high organic matter and moist soils.
Kudzu	Apply 8 oz/A as part of kudzu abatement program. Retreatment of any re-sprouting kudzu crowns following the initial treatment is necessary to fully control kudzu.
Unimproved Turf	1/2 to 2 oz/A depending on target weeds.

See product label for complete directions for use.



[nufarm.com/usivm](https://www.nufarm.com/usivm)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/usivm](https://www.nufarm.com/usivm) to download a full label.

©2024 Nufarm. Important: Always read and follow label directions. Clipper® is a trademark of Valent U.S.A. LLC. Helicopter image courtesy of [G. Keith Douce](#), University of Georgia, Bugwood.org



15-AG-0025-D