

This seed has been treated with SATIVA IM RTU (imidacloprid, metalaxyl, tebuconazole).

CAUTION **TREATED SEED**

DO NOT USE TREATED SEED FOR FEED, FOOD, OR OIL PURPOSE. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEED.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Seed Treated with SATIVA IM RTU is harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long pants, long-sleeved shirt and protective gloves when handling treated seeds.

Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Groundwater Advisory Statement: Metalaxyl is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds. Do not contaminate bodies of water when disposing of planting equipment wash water. Dispose of seed packaging or containers in accordance with local requirements.

USE RESTRICTIONS:

The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lbs.

DO NOT graze or feed livestock on treated areas for 45 days after planting.

REPLANTING / CROP ROTATION RESTRICTIONS:

In the event of a crop failure or harvest of a crop grown from SATIVA IMF MAX treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date SATIVA IM RTU treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

DISPOSAL:

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 031709
13754000

Sativa is a Registered Trademark of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

