

Truslate®

Get Tough on Weeds with Truslate®

Truslate® is a selective post-emergence herbicide that controls annual and perennial broadleaf weeds in wheat, barley, and oats not underseeded with a legume. Truslate is also registered for use in field corn, sweet corn and grasses grown for seed. It can be used to control tough weeds on Conservation Reserve Program (CRP) acreage and non-cropland. Difficult weeds like Canada thistle, kochia and wild buckwheat don't stand a chance against Truslate. With two powerful active ingredients in a convenient premix, Truslate offers a wide window of application for maximum flexibility and convenience.

KEY PERFORMANCE BENEFITS

- Controls more than 60 broadleaf weeds and suppresses more than 15
- Controls large kochia and herbicide-tolerant biotypes of kochia, plus Canada thistle and wild buckwheat
- Readily absorbed by roots and foliage, disrupting plant cell growth
- Rapidly translocated
- Wide window of application
- Can be tank-mixed with most other broadleaf and grass herbicides

EPA REG. NO.	71368-86
ACTIVE INGR.	clopyralid MEA salt (11.3%), fluroxypyr 1-methylheptyl ester (12.3%)
FORMULATION	emulsifiable concentrate
CHEM. FAMILY	pyridine
HRAC NO.	4
MODE OF ACT.	synthetic auxin growth regulator
REI	12 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2 x 2.5 gal, 30 gal, 250 gal tote, bulk



Grow a better tomorrow.

Truslate®

KEY USES

Corn; sweet, field
Grass seed
Permanent grasses in CRP non-cropland

KEY WEEDS CONTROLLED

(See product label for complete list)

Buckwheat, wild
Burdock, common
Chamomile, false
Chickweed
Clover; black medic, hop,
red, sweet, white
Cocklebur, common
Daisy, oxeye
Dandelion
Dock, curly
Groundsel, common
Hawksbeard; narrowleaf,
Hawkweed; orange, yellow
Hemp dogbane
Horseweed (*marestail*), field
Jimsonweed
Kochia
Lentils, volunteer
Lettuce, prickly
Mallow; common*, Venice
Morningglory
Nightshade; black, cutleaf,
eastern black, hairy
Pineappleweed*
Puncturevine
Purslane, common
Ragweed; common, giant
Sicklepod
Sowthistle; annual, perennial*, yellow
Thistle; common, bull, Canada, musk
Velvetleaf
Vetch
Wormwood, biennial

**suppression*

USE SUMMARY

Apply Truslate® to actively growing weeds keeping in mind that the optimum herbicidal activity occurs between 55°F to 75°F. Reduced efficacy may occur at temperatures below 45°F or above 85°F.

Do not apply through any type of irrigation system and spray drift to adjacent susceptible broadleaf plants or broadleaf crops.

Apply post-emergence from the three-leaf stage through flag leaf emergence. Apply before weeds exceed 4 inches in height or diameter. For Canada thistle control, apply after majority of basal leaves have emerged, but before budding.

The addition of MCPA ester, 2,4-D, dicamba or bromoxynil provides a broader spectrum of control of wild mustard, pigweed, lambsquarters and Russian thistle.

See label for complete directions for use.



NUFARMPPOINT.COM