

Clipper®



Tough. Fast. Gone.

Clipper® delivers fast and selective control of tough invasive and nuisance plants such as cabomba, watermeal, Eurasian watermilfoil, water lettuce, duckweed, giant salvinia and more. Plus, Clipper dissipates quickly from the water column and does not accumulate in sediment. And, Clipper offers reduced PPE requirements with 51% WDG formulation for easier application and handling.

CLIPPER DELIVERS UNBEATABLE AQUATIC WEED CONTROL

- Fast and selective control of tough invasive and nuisance plants
- Dissipates quickly from the water column and does not accumulate in sediment, giving it a favorable environmental profile
- 51% WDG means reduced personal protective equipment (PPE) requirements for ease of application, transport, and handling

EPA REG. NO.	59639-161
ACTIVE INGR.	flumioxazin (51%)
FORMULATION	water-dispersible granules
CHEM. FAMILY	N-phenylphthalimide
HRAC NO.	14
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	4 x 1 lb, 4 x 5 lb



Grow a better tomorrow.

Clipper®

KEY USES

For the management of undesirable aquatic vegetation in slow moving or quiescent waters

KEY WEEDS CONTROLLED

(See product label for complete list)

Alligator weed
Cabomba
Coontail
Curlyleaf pondweed
Duckweed
Eurasian watermilfoil
Fanwort
Filamentous algae
Frogs-bit
Green algae
Hydrilla
Pondweed
Sago pondweed
Variable-leaf watermilfoil
Water fern
Water lettuce
Water pennywort
Watermeal

APPLICATION RECOMMENDATIONS

Clipper® is a contact herbicide that quickly degrades in water. Good coverage and contact time is essential. Make applications with sufficient volume of water according to the label. When making applications to floating or emergent weeds, apply at 6 to 12 oz per acre and mix with an adjuvant approved for aquatic sites. For subsurface applications apply Clipper at 200 to 400 ppb in the water column. Please see label for information on the effects of high pH on plant response and efficacy.

PRECAUTIONS AND RESTRICTIONS

No restrictions for swimming, drinking, or fishing.

DO NOT apply to intertidal or estuarine areas.

In areas with dense weed vegetation only treat 1/2 the water body at one time and wait 10-14 days before treating the remaining area. DO NOT re-treat the same section of water within 28 days of application.

Treated water may not be used for irrigation purposes on food crops until at least five days after application.

Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the *Irrigation Restrictions Following Application* table below.

DO NOT use in water utilized for crawfish farming.

DO NOT re-treat the same section of water with Clipper more than 6 times per year.

DO NOT exceed 400 ppb of Clipper during any one application.

IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

APPLICATION METHOD	RATE	WATER DEPTH	TURF & LANDSCAPE ORNAMENTALS	GREENHOUSE & NURSERY ORNAMENTALS
Surface Spray	6-12 oz per surface acre	Greater than 3 feet	None	5 Days
		Less than 3 feet	12 Hours	5 Days
Subsurface	Less than 200 ppb	N/A	1 Day	5 Days
	200-300 ppb	N/A	2 Days	5 Days
	300-400 ppb	N/A	3 Days	5 Days

See label for complete application rates and recommendations.

FAST AND SELECTIVE CONTROL OF TOUGH INVASIVE AND NUISANCE PLANTS

Trial Work, M. Belland, Aquatic Control Technology, 2012 Application Date: May 17, 2012
Pond Location: Northwest Connecticut Target Plants: Curlyleaf pondweed (submersed) and filamentous, green algae (floating)



www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.
©2015 Nufarm. Important: Always read and follow label instructions. Clipper® is a trademark of Valent U.S.A. Corporation. 15-10-0021



Scan this code with your smartphone to visit CDMS and view the complete product label.