

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR VICTORY HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for this product and this supplement.




11901 S. AUSTIN AVE.
ALSIP, IL 60803

Victory™ Herbicide

EPA Reg. No. 71368-75

KEEP OUT OF REACH OF CHILDREN



CAUTION

This supplemental label expires May 1st, 2022 and must not be used or distributed after this date

**FOR USE ON SUNFLOWER HYBRIDS WITH THE DUPONT™
ExpressSun™ TRAIT FOR RESISTANCE TO VICTORY
HERBICIDE**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application. All applicable directions and precautions in the label booklet for this product must be followed.

Also refer to additional "INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, **DO NOT** USE THIS PRODUCT. Product must not be used in or near greenhouses.

Victory Herbicide is intended for application only to sunflower hybrids with the DuPont ExpressSun trait for resistance to Victory Herbicide, referred to below as "ExpressSun" sunflowers. Apply only on sunflower hybrids labeled "ExpressSun" and warranted by the seed supplier to have resistance to direct application of tribenuron, the active ingredient in Victory Herbicide. **DO NOT** apply this product to sunflower hybrids that lack resistance to Victory herbicide.

Apply Victory at a rate of 1/6 – 1/3 ounce (0.0078-0.0156 lbs ai) per acre. Add a Methylated seed oil (MSO) having at least 80% active ingredient strength at a rate of 1% v/v (1 gallon surfactant per 100 gal of spray solution).

Apply Victory to "ExpressSun" sunflowers any time from the 2 leaf stage of growth up to but not including the bud formation stage. Temporary crop yellowing may be observed shortly after application of Victory, especially when applied to crops growing under environmentally stressful conditions.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

POSTEMERGENCE SEQUENTIAL APPLICATIONS

Depending upon rainfall or other environmental conditions, annual weeds may have a second flush of germinating seedlings. To maximize control of such weeds, it may be necessary to apply this product a second time, 14 or more days after the first application. The combined rate of the postemergence applications cannot exceed 2/3 oz. of this product (0.031 lb/a.i.) per acre per year. For the control of annual grasses, apply a grass herbicide (i.e. ASSURE® II EPA Reg #352-541, Quizalofol-P-ethyl). Refer to the Quizalofol-P-ethyl product labeling for use rates, weed size, adjuvant selection, precautions, and restrictions. For maximum performance, apply Quizalofol-P-ethyl Herbicide at least one day before, or seven days after, the application of this product.

Sunflower (0512020)

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds, weeds that were beyond the maximum size at application, and/or weeds that emerge after an application.

- Cultivation up to 7 days before the postemergence application of this product may decrease weed control by pruning weed roots, placing the weeds under stress, and/or covering the weeds with soil and preventing coverage.
- To allow this product to fully control treated weeds, cultivation is not advised for 7 days after application.
- Optimum timing for cultivation is 7 – 14 days after a postemergence application.

PRECAUTIONS

- Application of Victory prior to the 2 leaf stage of “ExpressSun” sunflowers could result in crop injury.
- Avoid application to “ExpressSun” sunflower fields in which germination is uneven (i.e., some plants are outside the specified leaf stage for application), as crop injury may result.
- Application(s) to “ExpressSun” sunflowers that are, or have been, stressed by severe weather conditions, frost, abnormally hot or cold or wet or dry conditions, low fertility, drought, water saturated soil, disease and/or insect damage prior to application may result in crop injury. If the above stress conditions are expected to occur within 3 days after application of Victory to “ExpressSun” sunflowers, crop injury may also occur.

RESTRICTIONS

- **DO NOT** apply Victory to Sunflowers that are not “ExpressSun” sunflowers as severe crop injury or death of the plant will occur.
- **DO NOT** apply within 70 days of sunflower harvest.
- Re-treatment Interval is 14 days.
- Maximum single application rate is 1/3 ounce (0.0078 lb/a.i.) per acre. Maximum annual application rate is 2/3 oz. per acre (0.031 lb/a.i.) postemergence per year.
- **DO NOT** make more than 2 applications per calendar year.

Read the "Limit of Warranty and Liability" in the label booklet for Victory Herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Victory is a Trademark of Nufarm Americas Inc.

DuPont, ASSURE, and ExpressSun are registered trademarks or trademarks of its affiliates.