

ChampION⁺⁺™

FUNGICIDE / BACTERICIDE*

SUPERCHARGED COPPER FOR SUPERCHARGED CONTROL.

ChampION⁺⁺™ has the smallest, most consistent copper ions (particles) of any WG copper formulation. The result: better coverage and better disease control. And, ChampION⁺⁺ delivers all that with less environmental load than high dose copper products.

COMPETITIVE ADVANTAGES

SUPERIOR COVERAGE AND CONTROL

- Certified for use in organic crop production
- Small particle size (average = 1.2 micron) results in excellent coverage and disease protection
- Stable WG formula pours easily and disperses quickly
- High density copper product means lower use rates and smaller pack sizes
- Controls plant pathogenetic fungi and bacteria*
- OMRI Listed® for organic crop protection

*Non public health bacteria

AT-A-GLANCE INFO

EPA REG. NO.	55146-115
ACTIVE INGR.	copper hydroxide (46.1%)
FORMULATION	water-dispersible granule
CHEM. FAMILY	inorganic
GROUP NO.	M1
MODE OF ACT.	multi-site contact activity
REI	24 hours (greenhouse) 48 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	4 lb bag (10 per case), 10 lb bag (4 per case), 30 lb bag



MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Agri-Mycin® 50



Phostrol®

 **Nufarm**
Grow a better tomorrow

SUPERCHARGED COPPER FOR SUPERCHARGED CONTROL.

KEY USES

Citrus

Field crops; alfalfa, barley, corn, oats, peanut, potato, soybean*, sugarbeet, wheat

Grapes and berries; brambles, blueberry, cranberry, currant, gooseberry, kiwifruit, raspberry, strawberry

Hops

Tree crops; almond, apple, apricot, avocado, cherry, filbert, macadamia, nectarine, olive, peach, pear, pecan, pistachio, plum, prune, quince, walnut

Tropical crops; banana, cacao, coffee, guava, litchi, mamey sapote, mango, papaya, passion fruit, plantain, sugar apple

Vegetables; bean, beet, carrot, celery, celeriac, crucifers, cucurbits, eggplant, endive, escarole, garlic, leek, lettuce**, okra, onion, pea, pepper, spinach, tomato, watercress

*Not for use in California

**Not for use in California and Arizona

KEY DISEASES CONTROLLED

(See product label for complete list)

Algal leaf spot	Crown rot
Alternaria	Downy mildew
brownspot	Fire blight
Anthracnose	Gummy stem
Bacterial blast	blight
Bacterial blight	Leaf curl
Bacterial spot	Leaf rust
Black pitting	Leaf spot
Black rot	Phytophthora
Botrytis blight	brown rot
Brown spot	Phytophthora
Cane spot	foot rot
Cercospora	Powdery mildew
leaf blight	Pseudomonas
Cercospora	blight
leaf spot	Scab
Citrus canker	Spanish moss
(suppression)	Walnut blight
Collar rot	Yellow rust

USE SUMMARY

RATES: PER APPLICATION

Citrus.....	0.5 lbs/A
Field crops	0.5-2.0 lbs/A
Grapes and berries	0.75-4.25 lbs/A
Hops.....	0.75-1.5 lbs/A
Tree crops.....	0.5-10.5 lbs/A
Tropical crops.....	0.75-7.75 lbs/A
Vegetable crops.....	0.5-3.5 lbs/A

APPLYING SPRAY MIXTURE: The amount of ChampION⁺⁺™ applied per acre in concentrate and aerial sprays must be the same as the amount applied per acre in dilute sprays. The required amount of ChampION⁺⁺ must be mixed with enough water to thoroughly cover the crop with spray mixture and is to be applied to the point of runoff for best performance. The volume of water needed depends upon factors such as the spray equipment used, foliage density, size of the crop being sprayed, etc.

CHEMIGATION APPLICATION: Apply ChampION⁺⁺ only through sprinkler irrigation systems including center pivot, motorized lateral move, end tow, traveler, big gun, plastic solid set, or plastic hand move; DO NOT apply ChampION⁺⁺ through any other type of irrigation system. In California, do not apply in systems which contain aluminum parts or components.

See label for complete directions for use.

APPLICATION PRECAUTIONS

DO NOT apply ChampION⁺⁺ in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of this product resulting in possible phytotoxicity or loss of effectiveness.

DO NOT tank mix with other pesticides or additives unless the mixture has been determined to be compatible and non-injurious to the crop under your conditions of use.

See label for complete application precautions.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2020 Nufarm. Important: Always read and follow label instructions. Agri-Mycin®, ChampION⁺⁺™, Grapple™ and Phostrol® are trademarks of Nufarm. OMRI Listed® is a trademark of Organic Materials Review Institute.

