

Abamex™

INSECTICIDE

LOW VOC,
WATER-BASED
FORMULA

TARGET AND MANAGE TOUGH MITES.

Historically, abamectin has provided cotton growers with a powerful tool to combat difficult-to-control spider mites and other mite species. Abamex™ packages that power in a low-volatile, water-based formulation. Abamex starts working quickly to affect mites within hours of exposure. Its translaminar activity means it protects the entire leaf, inside and out. And, selective chemistry makes it a useful rotation partner for resistance management programs.

COMPETITIVE ADVANTAGES

CONVENIENT CONTROL

- Effective management of two-spotted spider mites and other difficult-to-control mite species
- Translaminar leaf penetration
- Flexible application timing in an easy-to-use, low-volatility liquid formulation
- Rainfast within hours of application

AT-A-GLANCE INFO

EPA REG. NO.	228-734
ACTIVE INGR.	abamectin (2.0%)
FORMULATION	microemulsion concentrate (MEC)
CHEM. FAMILY	avermectin
IRAC NO.	6
MODE OF ACT.	chloride channel modulator
REI	12 hours
SIGNAL WORD	caution
PACKAGE SIZE	1 gal container (4 per case)

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Goal® 2XL | Trunemco™ | Panther SC

 **Nufarm**
Grow a better tomorrow



TARGET AND MANAGE TOUGH MITES.

KEY USES

Cotton

KEY INSECTS CONTROLLED

(See product label for complete list)

Carmine spider mite
Pacific spider mite
Strawberry spider mite
Two-spotted spider mite

USE SUMMARY

COTTON PHI: 20 days

EARLY-SEASON: 4.0–6.0 fl oz/A

Apply this product when mites first appear.

Rate will depend on the cotton plant size and thickness of foliage. Use the 4.0–6.0 fl oz/A rate only early in season and for cotton plants 10 inches or less in height.

Applications can be made by air or ground using a minimum of 5 gallons of water/A.

West of the Rockies: DO NOT apply less than 4.0 fl oz/A and can only be applied by ground equipment.

MID-SEASON/LAY-BY: 8.0–16.0 fl oz/A

Apply this product when mites first appear.

Rate will depend on the cotton plant size and thickness of foliage.

Applications can be made by air or ground using a minimum of 5 gallons of water/A.

ABAMEX™ CONTROL OF TWO-SPOTTED SPIDER MITES – CA 2013 COTTON

	PRE-TREAT	3 DAT	7 DAT	14 DAT	21 DAT
UTC (UNTREATED)	1.5	2.8	2.3	0.5	0
ABAMEX 4 FL OZ	1.0	0.8	0.8	0	0
ABAMEX 8 FL OZ	0.5	0.5	0.5	0	0

NIS at 1% v/v added to all treatments

Counts represent number of two-spotted spider mites per leaf
Trial conducted on cotton

Abamectin treatments were significantly different from the UTC but not each other at 3 and 7 DAT

See product label for complete directions for use.

APPLICATION RESTRICTIONS

DO NOT apply more than 16 fl oz/A per application.

DO NOT make more than 2 applications per season with a 21-day interval between applications.

DO NOT exceed 32 fl oz/A per growing season.

Pre-harvest Interval: 20 days

DO NOT graze livestock in treated fields or feed cotton foliage to livestock.

See product label for complete directions for use.



nufarm.com/uscrop

Abamex™ is a Restricted-Use Pesticide. Always read and follow directions. For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2020 Nufarm. Important: Always read and follow label instructions. Abamex™ and Panther® are trademarks of Nufarm. Trunemco™ is a trademark of BASF and exclusively marketed by Nufarm. Goal® is a registered trademark of Nutrilchem Co. Ltd. 15-AG-0100-B-Cotton

