

# GoalTender®

SELECTIVE HERBICIDE

## STAY UNDEFEATED WITH GOALTENDER®

GoalTender® herbicide is a water-based formulation that holds well to soil and performs at lower rates with less environmental load. It delivers burndown of existing broadleaf weeds and provides long-lasting pre- and post-emergence activity on dozens of broadleaf weeds and grasses, making it ideal for residual herbicide programs in tree nut, tree fruit, vine and vegetable crops. GoalTender is an excellent choice for VOC requirements compared to higher VOC formulations.

### COMPETITIVE ADVANTAGES

#### TRUSTED BRAND

- Wide range of labeled crop and application timings that are tried and tested
- Gold standard in tree nut, fruit and vine, providing residual herbicide programs with proven staying power, up to 8 months

#### BROAD SPECTRUM

- As pre-emergence, GoalTender forms a chemical barrier on the soil surface and affects weeds through direct contact at emergence
- GoalTender has the added benefit of post-emergent activity on many actively growing labeled weeds
- Holds well to soil with low volatility
- Excellent tank mix partner
- Water-based formulation allows for more springtime applications with less off-target potential, greater crop safety and application flexibility

#### LONG LASTING

- Offers residual control in tree nuts, cotton, grapes, tree fruit, and many vegetables to keep weeds clear for months

#### LOW LEACHING OR VOLATILIZATION

- Best pre-emerge control is achieved when irrigation or rainfall (at least .25 inches) occurs within 3–4 weeks after application

### AT-A-GLANCE INFO

EPA REG. NO. 92894-3-71368

ACTIVE INGR. oxyfluorfen (41%)  
4 lbs/gal

FORMULATION liquid

CHEM. FAMILY diphenyl ethers

GROUP NO. 14

MODE OF ACT. PPO inhibitor

REI 24 hours,  
48 hours (onions, garlic,  
and horseradish)

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 1 gal (4 per case)

MORE PRODUCTS EXCLUSIVELY FROM NUFARM



# GoalTender®

## STAY UNDEFEATED WITH GOALTENDER®

### KEY USES

Many crops including almond\*, apple, cherry, cotton, grape\*\*, citrus, olive, onion (direct seeded, transplanted, grown for seed), pear, pecan, pistachio\*, and walnut\*

\* includes non-dormant application in CA and AZ only

\*\* includes non-dormant application in CA only

### KEY WEEDS CONTROLLED

(See product label for complete list)

#### COMMON BROADLEAF WEEDS AND GRASSES:

Barnyardgrass  
Crabgrass, large  
Cutleaf evening primrose\*  
Fescue  
Foxtail, giant  
Goosegrass  
Groundsel, common  
Henbit  
Kochia  
Lambsquarters  
Mallow, little  
Morningglory species, annual  
Nightshade, American black and black prickly sida  
Oats, wild  
Pigweed, common and redroot  
Pennsylvania smartweed  
Shepherd's-purse  
Sowthistle  
Spurge, prostrate and spotted  
Velvetleaf

\*highest rate and/or multiple applications may be required for acceptable control

### USE SUMMARY

RATE (PT/A)	RECOMMENDATION
<b>DORMANT TREE FRUIT, NUT, AND VINE CROPS</b>	
2.5-3 (pre-emerge broadcast)	Apply a min. of 20 gal of water/A. Use higher spray volumes in high densities of emerged weeds or heavy trash. Sprays should be directed to the soil and the base of dormant trees or vines.
2.5-4 (pre-emerge banded)	
1-3 (post-emerge broadcast)	Apply in a spray volume of 40 or more gal/A. For best control, apply when weeds are at seedling stage. Lower rate (1 pt/A) is specified for control up to 4-leaf stage. Higher rates (up to 3 pt/A) may be used for weeds up to 6-leaf stage. Applications beyond the 6-leaf stage may result in partial control.
1-4 (post-emerge banded)	

**Precautions:** Apply to only healthy growing trees or vines. Avoid direct plant contact. Direct spray toward the base of tree or vines unless specific use recommendations allow over-the-top application.

**Restrictions:** Unless otherwise specified, do not apply between bud swell and completion of final harvest or when fruit/nuts are present.

In AZ and CA, may apply following final harvest up to Feb 15 (Feb 1 in Coachella Valley, CA). Applications after these dates, but prior to bud swell, may result in significant crop injury.

For banded applications, up to 4 pt/A per use season may be applied within the treated band. Do not apply more than a max of 3 pt/A per use season on a broadcast basis.

Do not apply to grapes or kiwi established less than 3 years unless vines are on a trellis wire a minimum of 3 feet above the soil surface.

Do not apply to grapes or kiwi that are not staked or trellised unless vines are free standing. Maximum total application rate per year is 1.5 lbs ai/A.

#### NON-DORMANT GRAPES (CALIFORNIA ONLY)

1 (pre-emerge)	Apply as a directed spray in min. spray volume of 20 gal/A or through low-vol sprinkler or drip irrigation. Add 1 qt 80% active nonionic surfactant/100 gal spray for post-emerge direct spray.
0.5-1 (post-emerge)	

**Precautions:** GoalTender may result in injury to non-dormant grapes including necrosis, reddening, cupping or crinkling of leaves. The plant will continue to grow normally.

Grapes may exhibit some small blemishes (spots or flicks) on the fruit. GoalTender is phytotoxic to plant foliage. Avoid drift to all other crops and non-target areas.

**Restrictions:** The total amount applied during one season cannot exceed 3 pt/A.

Do not apply within 14 days of harvest.

Do not initiate application in non-dormant grapes until the completion of bloom period.

Do not apply to grapes established less than 3 years unless vines are either on a trellis wire a minimum of 3 feet above the soil surface, or protected by grow tubes.

Apply only by ground application equipment or low-volume sprinkler or drip irrigation.

Apply as a non-dormant application to wine grapes or raisin grapes only.

#### NON-DORMANT PISTACHIO, WALNUT, AND ALMOND (CALIFORNIA AND ARIZONA ONLY)

2.5-3 (pre-emerge)	For residual weed control of listed weeds.
0.5-1 (post-emerge suppression)	Apply to weeds < 4" in height. Repeat applications may be required.
1-3 (post-emerge cleanup)	For cleanup or pre-harvest applications, apply to weeds < 4" in height. Applications beyond the 4" stage may result in partial control.

**Precautions:** Direct spray toward the base of trees. Avoid direct contact with foliage or nuts. Apply only to healthy growing trees.

**Restrictions:** As a non-dormant treatment, GoalTender can only be applied to pistachio plantings between May and 7 days prior to harvest, to almond plantings between April 1 and September 30 and to walnut plantings between May 1 and September 30.

Do not apply within 7 days of pistachio harvest, within 30 days of almond harvest or within 7 days of walnut harvest.

Do not apply more than 3 pt/A during the non-dormant season.

Maximum total application rate per year is 1.5 lbs ai/A.

See label for complete directions for use.



[nufarm.com/uscrop](http://nufarm.com/uscrop)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/uscrop](http://www.nufarm.com/uscrop) to download a full label.

©2021 Nufarm. Important: Always read and follow label instructions. Cheeta®®, Grapple™ and Tuscany® are trademarks of Nufarm. GoalTender® is a registered trademark of Nutrichem Co. Ltd.

