



ACHIEVE OPTIMUM RESULTS.

BAHIA GRASS, BERMUDAGRASS AND ZOYSIAGRASS

For optimum and efficient performance apply this product at the rate of 2-3 pts in 20-240 gals of water/A (0.73-1.10 fl oz in 0.5-5.5 gals of water/1,000 sq ft) to control weeds when temperatures are below 80°F. Apply at 1-2 pts/A between 80 and 90°F. Avoid broadcast applications when air temperature exceeds 90°F. When air temperature exceeds 90°F limit application to spot treatment only. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur. Avoid double coverage resulting from overlap unless temporary turf injury can be tolerated.

CARPETGRASS, CENTIPEDEGRASS AND ST. AUGUSTINEGRASS (EXCEPT FLORATAM AND BITTERBLUE)

For optimum and efficient performance apply this product at the rate of 1-2 pts in 20-240 gals of water/A (0.37-0.73 fl oz in 0.5-5.5 gals of water/1,000 sq ft) to control weeds. Apply at 1 pt/A between 80°F and 90°F. Avoid broadcast applications when air temperature exceeds 90°F. When air temperature exceeds 80°F limit application to spot treatment only. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur. Avoid double coverage resulting from overlap unless temporary turf injury can be tolerated.

FOR OPTIMUM RESULTS

(1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1-2 days before and following application; (3) reseed no sooner than 3-4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass or Bitterblue. The application of this herbicide to grass seedlings is recommended after the second mowing. The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3-4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant zoysiagrass and dormant bahiagrass are suggested.

NOTE: For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc. For residential turf and golf course uses, do not apply more than 1.5 lb ai/A (3 pts product)/year. Do not apply more than 2 applications/year with a minimum re-treatment interval of 21 days.

Do not apply this product in a way that will contact workers, any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not use on Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Aerial application is prohibited.