

# 26/36<sup>®</sup>

FUNGICIDE

## DUAL MODE OF ACTION FOR EFFECTIVE DISEASE CONTROL.

Incorporating 26/36<sup>®</sup> Fungicide into an integrated disease management program provides effective control of key diseases of turfgrass and ornamentals.






### COMPETITIVE ADVANTAGES

#### EASY-TO-USE FORMULA

- Easy-to-use, premixed formulation
- Convenient
- Broad-spectrum disease control for turf
- Tank-mix compatible with most commonly used pesticides

### AT-A-GLANCE INFO

EPA REG. NO.	228-630
ACTIVE INGR.	iprodione (19.65%), thiophanate-methyl (19.65%)
FORMULATION	flowable
CHEM. FAMILY	dicarboxamide, benzimidazole
FRAC NO.	1 and 2
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2 x 2.5 gal

-  COMMERCIAL
-  TURF
-  LANDSCAPE ORNAMENTALS
-  GOLF TURF
-  SOD FARMS

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

**3336<sup>®</sup>** | **Arena<sup>®</sup>**  
INSECTICIDE | **Celero<sup>®</sup>**  
HERBICIDE

 **Nufarm**  
Grow a better tomorrow

# 26/36®

## DUAL MODE OF ACTION FOR EFFECTIVE DISEASE CONTROL.

### KEY USES

Golf courses  
Institutional areas  
Ornamentals  
Sod farms  
Turfgrass

### KEY DISEASES CONTROLLED

(See product label for complete list)

Anthrachnose  
Ascochyta leaf blight  
Brown patch  
Copper spot  
Corticium  
Dollar spot  
Fusarium patch  
Gray snow mold  
Leaf spot  
Pink snow mold  
Red thread  
Zoysia patch

### USE SUMMARY

26/36® Fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. It provides effective, broad spectrum turf disease control and also serves as a useful tank mixture in the resistance management program required for other resistant sensitive fungicides.

When used in conjunction with good turf management practices, 26/36 Fungicide is effective in controlling the following diseases: dollar spot, brown patch, anthracnose, leaf spots such as Helminthosporium leaf spot caused by Drechslera spp. pathogens, corticium red thread, Fusarium patch, zoysia patch, ascochyta leaf blight and copper spot.

Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

[See label for complete directions for use.](#)

### APPLICATION PRECAUTIONS AND RESTRICTIONS

DO NOT exceed a total of 14.5 fl oz of product per 1000 sq ft per year.

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. When disease pressure is low to moderate, the lower rates and longer intervals are recommended.

DO NOT mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred.

DO NOT mix with any sticker, extender, or wetting agent.

DO NOT graze animals on treated turf.

DO NOT feed clippings from treated turf to livestock or poultry.

[See label for complete application precautions and restrictions.](#)



[nufarminsider.com](http://nufarminsider.com)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/us](http://www.nufarm.com/us) to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. 26/36® and 3336® are trademarks of Nufarm. Arena® and Celero® are trademarks of Sumitomo Chemical Company, Ltd.



15-TO-0029-D