

Arena[®] 50 WDG

INSECTICIDE

A BETTER VALUE THAT LASTS. AND LASTS. AND LASTS.™

Don't settle for insecticides that give up on control early. Get the best control and the longest-lasting residual out there with Arena[®] Insecticide. Arena works as both a preventive and a curative, for outstanding control of chinch bugs, white grubs and a broad spectrum of other turf and ornamental pests with a single application.

COMPETITIVE ADVANTAGES

PROVEN RESIDUAL ACTIVITY FOR LASTING CONTROL

- Controls pyrethroid-resistant chinch bug, without increasing the level of pyrethroid resistance
- Controls white grubs, including Japanese beetle and masked chafers, plus other thatch- and surface-feeding insects
- Superior grub control including curative grub control that helps keep away foraging animals
- Excellent results when applied prior to third instar
- Does not require immediate watering-in after application
- Also available as a 0.25 granule formulation for convenient broadcast spread application

AT-A-GLANCE INFO

EPA REG. NO.	59639-152
ACTIVE INGR.	clothianidin (50%)
FORMULATION	water-dispersible granules
CHEM. FAMILY	neonicotinoid
IRAC NO.	4A
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	4 x 40 oz bottle

 RESIDENTIAL

 COMMERCIAL

 TURF

 GOLF TURF

 LANDSCAPE ORNAMENTALS

 SOD FARMS



Arena® 50 WDG

INSECTICIDE

A BETTER VALUE THAT LASTS. AND LASTS. AND LASTS.™

KEY USES

Commercial and residential turf and landscape ornamentals
Golf courses
Interior plantscapes
Non-bearing fruit and nut trees
Recreational areas
Residential apple and pear trees
Sod farms

KEY PESTS CONTROLLED

(See product label for complete list)

Aphids
Armyworms
Asiatic garden beetle
Azalea lace bug
Billbugs
Black turfgrass atatenius
Caterpillars
Cutworms
European chafer
European crane fly
Florida wax scale
Glassy-winged sharpshooter
Green June beetle
Japanese beetle
Leafhoppers
Leafminers
Mealybugs
Mole crickets (suppression)
Northern masked chafer
Nuisance ants
Oriental beetle
Phyllophaga spp. (June beetles)
Pyrethroid-resistant chinch bug
Root weevils
Scales
Sod webworms
Southern masked chafer
Spittle bug
White grubs
Whiteflies

USE SUMMARY

Use rates from 4.5 to 12.8 oz/A. (Rate depends on pest and time of application.) 9.6 oz for white grub curative and southern chinch bug.

See label for complete directions for use.

PRECAUTIONS AND RESTRICTIONS

DO NOT use a foliar application of clothianidin following a soil application of clothianidin.

Regardless of formulation or type of application (soil or foliar), DO NOT apply more than a total of 0.4 lb clothianidin per acre per season for turf, landscape, ornamentals, and non-bearing fruits and nuts.


If the maximum seasonal limit of clothianidin as mentioned above has been applied and pest populations require additional treatments, use another registered pesticide that is not in the neonicotinoid class of chemistry.

DO NOT apply by air.

See label for complete application rates and recommendations.

TURF INJURY FROM SOUTHERN CHINCH BUG

Source: Dr. Eileen Buss, University of Florida, 2010



nufarminsider.com

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. Arena® is a trademark of Sumitomo Chemical Company, Ltd. A better Value that lasts. And lasts. And lasts.™ and Aloft® are trademarks of Valent U.S.A. LLC. Talstar® is a trademark of FMC Corp.



15-TO-0008-B