

Triplet® Low Odor



Powerful Weed Control Without the Powerful Smell.

Triplet® Low Odor is a liquid post-emergent broadleaf herbicide that uses an innovative and unique formulation of 2,4-D. This powerful combination herbicide selectively controls broadleaf weeds in turfgrass, without the initial strong odor or lingering chemical smell. Stubborn weeds like dandelion, clover, chickweed, henbit and plantains don't stand a chance against the three dynamic herbicides in Triplet Low Odor.

BROAD-SPECTRUM CONTROL WITH NO RESIDUAL SMELL

- Efficacious, broad-spectrum weed control with significantly reduced odor and no residual smell
- Offers the power of three proven herbicides including a unique TIPA 2,4-D
- Proprietary surfactant system that enhances weed penetration and outperforms traditional three-way products
- Excellent control of a variety of broadleaf weeds
- Ideal for cool- or warm-season turf
- Labeled for sod farm usage
- Tank-mix compatible with most liquid fertilizers and iron*
*Compatibility test required

EPA REG. NO.	228-409
ACTIVE INGR.	2,4-D TIPA (47.33%), mecoprop-p (8.17%), dicamba (2.30%)
FORMULATION	liquid
CHEM. FAMILY	phenoxyaliphatic acid, benzoic acid
HRAC NO.	4
SIGNAL WORD	danger
RESTRICTED	no
PACKAGE SIZE	2 x 2.5 gal, 30 gal, 55 gal



RESIDENTIAL



COMMERCIAL



TURF



GOLF TURF



POST-EMERGE



SOD FARMS



Nufarm

Grow a better tomorrow.

Triplet® Low Odor

KEY USES

Athletic and sports fields
 Cemeteries
 Golf courses (*fairways, aprons, tees and roughs*)
 Non-turf areas (*roadsides and rights-of-ways*)
 Ornamental turf lawns (*residential, industrial and institutional*)
 Parks
 Sod farms

KEY WEEDS CONTROLLED

(See product label for complete list)

Bedstraw
 Black medic
 Buckhorn
 Burdock
 Carrot, wild
 Chicory
 Chickweed
 Clover
 Dandelion
 Dock
 Garlic, wild
 Ground ivy
 Healall
 Henbit
 Knotweed
 Lambsquarters
 Lespedeza
 Lettuce, wild
 Mallow
 Morningglory
 Onion, wild
 Peppergrass
 Pigweed
 Plantain
 Poison ivy
 Poison oak
 Purslane
 Ragweed
 Sheep sorrel
 Shepherd's-purse
 Speedwell
 Spurge
 Yarrow

USE SUMMARY

ORNAMENTAL LAWNS & TURF APPLICATION RATES

SITE	USE RATE (fluid ounces)	SPRAY VOLUME (gallons)	USE RATE (pints)	SPRAY VOLUME (gallons)
	PER 1000 SQ. FT.		PER ACRE	
Bahiagrass, bluegrass, common bermudagrass, fescue, ryegrass, zoysiagrass	1.1 to 1.5	0.5 to 5	3 to 4	5 to 220

Use Directions: For Professional Lawn Maintenance: Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

If Bermudagrass is Dormant: Up to 4 pints per acre may be used. However, some hybrid bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.

St. Augustinegrass and centipedegrass	0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200
---------------------------------------	-------------------------	--------	---------------------	-----------

Use Directions: Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy. Some temporary discoloration may occur on warm-season grasses.

Bentgrass (putting and bowling greens)	0.67 Maximum	3.3	1.8 Maximum	145
--	-----------------	-----	----------------	-----

Use Directions: Apply on closely mowed bentgrass, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

Note: Care should be taken to avoid overdosing bentgrass, St. Augustinegrass and centipedegrass or injury may result. Large volumes of spray water (i.e. 0.67 fluid ounce in 3.3 gallons of water per 1,000 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray is from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

SOD FARM APPLICATION RATES

SITE	USE RATE (pints)	SPRAY VOLUME (gallons per acre)
Bluegrass, fescue, ryegrass	3 to 4	5 to 175
Creeping bentgrass	1.8	
Bahiagrass, common and hybrid bermudagrass, St. Augustinegrass and zoysiagrass	2 to 2.5	

Use Directions: Apply to actively growing broadleaf weeds. Application to grass seedlings is recommended after the second mowing.

NON-TURF AREAS APPLICATION RATES

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 7 pints per 43,560 sq ft. Ensure adequate water to thoroughly saturate all weeds.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 1 to 1.5 gallons per 43,560 sq ft.

See label for complete directions for use.

