

Tourney® FUNGICIDE

TURF & ORNAMENTAL DISEASE CONTROL

Tourney® Fungicide helps you defeat the toughest diseases in turfgrass and ornamental trees and shrubs single-handedly, saving time and money.

TOURNEY DELIVERS

- Control of more than 16 tough turf and ornamental diseases – including apple scab, brown patch, powdery mildew, rust, anthracnose, leaf spots, conifer blights and more – for up to 28 days
- Low use rates that allow you to do more with less, saving you money
- Application flexibility – available as a foliar spray or soil drench for root and crown diseases


APPLICATION INFORMATION

DISEASES CONTROLLED		RATE (OZ/100 GALLONS WATER)
<ul style="list-style-type: none"> • Anthracnose • Apple Scab • Botrytis • Blossom Rot/Brown Rot • Brown Patch • Conifer Blights 	<ul style="list-style-type: none"> • Damping Off/Stem & Root Rots • Eastern Filbert Blight • Leaf Spots • Powdery Mildew • Rust • Shot Hole 	<p>2 to 4 oz on a 14- to 28-day interval.</p> <p>Apply as a foliar spray when targeting foliar diseases or as a drench when targeting crown and/or root diseases.</p> <p>100 gallons of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.</p>

TOURNEY DELIVERS OUTSTANDING LILAC POWDERY MILDEW

Tough Disease Control

Tourney performs well even in high disease severity situations. The chart below demonstrates how Tourney delivers outstanding powdery mildew control for plants like lilac either solo or combined with another fungicide.



B. Rao, Davey Tree. Application intervals: Tourney (2 oz/A) - 14 days; others - 28 days. Applications initiated June 27.



Nufarm

Grow a better tomorrow.

Tourney® FUNGICIDE

One of the most common diseases on crabapple and non-bearing apple trees for nurseries and landscape settings in cool, humid regions is apple scab, caused by *Venturia inaequalis*. Unsightly leaf spots are the most visible and offensive symptoms of apple scab, but petioles, blossoms and fruit are also susceptible to infection. Extensive leaf infections result in leaf defoliation. Successive years of leaf defoliation weaken the tree, leading to lower quality blooms in subsequent years and increased sensitivity to damage from cold temperatures.


Tourney® provides excellent control of apple scab, even when applied on a 28-day interval.

TO BEST CONTROL APPLE SCAB, APPLY TOURNEY

- At 2 to 4 oz product per 100 gallons water
- As a full coverage, foliar spray to the point of run-off
- On a 14- to 28-day interval

APPLE SCAB CONTROL, CRABAPPLE (*MALUS SP.*) TREES

Apple scab can be an especially tough disease and Tourney delivers best-in-class control even on 28-day spray intervals. The chart below demonstrates the superior apple scab control of Tourney vs. the industry standard.



B. Rao, Davey Tree. Treatments were applied 23 April, 5 May (14 day interval only), & 21 May. Natural infections of *Venturia inaequalis* were rated. Rates are ounces per 100 gallons water.



www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.
©2015 Nufarm. Important: Always read and follow label instructions. Tourney® is a trademark of Valent U.S.A. Corporation. Banner MAXX® is a trademark of Syngenta Group Company. 3336® is a trademark of Nufarm. 15-TO-0079