

TriStar[®] 8.5 SL

INSECTICIDE

THE CHOICE FOR BASAL BARK TREATMENT.

Apply TriStar[®] 8.5 SL as a basal bark application to combat many woody ornamental pests, including borers, scales, and adelgids, in hard-to-spray areas. This easy-to-use water-based formula works through translaminar activity to distribute treatment throughout the plant – providing lasting insect protection to old and new growth.

COMPETITIVE ADVANTAGES

QUICK KNOCKDOWN AND LASTING RESIDUAL CONTROL

- For use on ornamental or non-fruit bearing fruit and nut trees
- Controls borers, scale insects, and many other insects
- For low pressure small volume sprayers mix 12.5-25.3 fl oz of TriStar per gallon of water with an organo-silicate adjuvant and apply 3-4 fl oz of spray mixture per inch DBH
- Depending on insect target, treatments begin at bud break through full leaf expansion in early to mid-spring

SOLUTION TIP: WHEN TO CHOOSE WHICH PRODUCT

1. USE TRISTAR[®] WHEN POLLINATOR IMPACT IS A CONCERN.

TriStar (acetamiprid) is not affected by current US regulatory actions or previous EU regulatory actions. In fact, TriStar is an excellent choice for landscape pros looking to control insect pests while reducing the risk to pollinator species.

See the TriStar Pollinator Sheet at Nufarm.com/USTurf/tristar

2. USE SAFARI[®] WHEN EMERALD ASH BORER CONTROL IS NEEDED.

Safari is a new generation, super-systemic insecticide that provides control of many insect pests including effective control of emerald ash borer (EAB), a devastating invasive pest that has killed tens of millions of ash trees. Safari controls EAB when applied as a soil drench, soil injection or basal trunk spray.

See the more about EAB control at Nufarm.com/USTurf/safari

AT-A-GLANCE INFO

EPA REG. NO. 8033-106-1001

ACTIVE INGR. acetamiprid (8.5% by wt.)

FORMULATION soluble liquid

CHEM. FAMILY neonicotinoid

IRAC NO. 4A

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 1 qt (4 per case),
1 gal (4 per case)



RESIDENTIAL



COMMERCIAL



LANDSCAPE ORNAMENTAL

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

4-Speed XT

Arena
INSECTICIDE

Q-Ball
HERBICIDE

 **Nufarm**
Grow a better tomorrow

THE CHOICE FOR BASAL BARK TREATMENT.

KEY USES

Annual and perennial
flowering plants
Ornamental plants
Shrubs
Vines

KEY INSECTS CONTROLLED

(See product label for complete list)

Adelgids
Aphids
Citrus thrips
Crane fly larvae
European pine sawflies
Hard and soft scales
Leaf eating beetles
Leafhoppers
Leafminers
Mealybugs
Plant bugs
Psyllids
Strawberry weevils
Thrips
Whiteflies

USE SUMMARY

FOR BASAL BARK TREATMENT OF TREES (ORNAMENTAL OR NON-BEARING FRUIT AND NUT): TriStar[®] 8.5 SL Insecticide may be applied as an external basal bark application to ornamental and non-bearing fruit and nut trees for control of borers (such as Flathead Apple Borer, but excluding Emerald Ash Borer or Asian Longhorn Beetle), scale insects (such as Calico Scale, Gloomy Scale, Azalea Bark Scale) and Hemlock Woolly Adelgids in hard to spray or environmentally sensitive areas. This application involves wetting the bark of the tree starting from a height of approximately 8 feet downward to the exposed root flair with a directed spray to completely wet the application area. Depending on the specific bark characteristics, use 3-4 fluid ounces of spray per inch Diameter Breast Height (DBH at 4.5 feet) of the intended target.

See product label for complete directions for use.

APPLICATION INSTRUCTIONS

Depending on insect target, treatments may begin at bud break through full leaf expansion in early to mid-spring. Consult your local extension service recommendations for the target pest. Make applications as required for preventive or curative management of pest.

Using a low pressure (10-25 PSI), small volume handcan or backpack sprayer, mix 12.5-25.3 fl oz of TriStar 8.5 SL per gallon of water with an organo-silicate adjuvant according to the adjuvant product instructions. Apply as a full coverage spray starting at the top of the application zone and working downward to the root flair. One gallon should treat approximately 36-42 total inches of treatment DBH depending on bark surface.

DO NOT apply to wet bark, or during rainfall, or if rain is expected within 12 hours after application.

DO NOT apply as a drench to the soil.

See product label for complete application instructions.

