

Curator™

Protect Your Masterpiece

Pythium blight, damping off and root rot can strike suddenly during hot, humid weather, especially in the wettest areas. Curator™ Fungicide is your best defense against these devastating diseases with a unique combination of active ingredients to enhance your pythium management program as a rotational partner with multiple modes of action. This cost-effective treatment, backed by the reliability of dual-active ingredients, saves both time and money.

COMPETITIVE ADVANTAGES

- Two active ingredients, propamocarb and mefenoxam, enhances resistance management and overall performance.
- Effective prevention and control against pythium blight, damping off and root rot diseases.
- Fast-acting activity provides immediate disease protection by stopping germinating spores and halting mycelium growth on contact.
- Systemic activity offers rapid control and lasting protection with root and foliar absorption, and upward mobility.
- Highly effective against pythium crown and foliar blight control in cool-season turfgrass with two modes of action for resistance management.
- A first line of defense for crown and foliar pythium in warm-season turfgrass, fitting seamlessly into pythium management programs.
- Effective control of downy mildew in St. Augustinegrass and yellow tuft (downy mildew) in *Poa annua* (bluegrass).

AT-A-GLANCE INFO

EPA REG. NO.	228-771
ACTIVE INGR.	propamocarb (3.6 lb/gal), mefenoxam (0.36 lb/gal)
FORMULATION	micro-emulsion liquid
CHEM. FAMILY	carbamate, acylalanines
FRAC NO.	28 & 4
SIGNAL WORD	caution
PACKAGE SIZE	2 x 2.5 gal

Curator™

KEY USES

Seedling and established cool- and warm-season turf, including overseeded turfgrass for: golf course, athletic fields, lawns around residential, institutional, public, commercial and industrial buildings; parks, recreational areas, sod farms

KEY DISEASES CONTROLLED

(See product label for complete list)

Damping off
Pythium blight
Pythium root rot

USE SUMMARY

For best efficacy apply as a preventative treatment at 3-6 fl oz/1000 sqft. Irrigation (1/8-1/4 inch) or rainfall, is required after application.

Under severe disease conditions, use the highest labeled rate on a 7-14 day schedule.

ESTABLISHED TURF

Apply when conditions favor disease development. Favorable conditions include high temperatures and high humidity.

OVERSEEDDED TURF

Apply after seed germination to prevent pythium damping off.

PYTHIUM BLIGHT CONTROL

Control of Pythium Blight on Perennial Ryegrass (14-day Interval)

2022 RUTGERS TURF GRASS FIELD DAY



Curator 3 fl oz



Curator 6 fl oz



Check

2023 RUTGERS TURF GRASS FIELD DAY



Curator 6 fl oz



Segway® 0.6 fl oz



Serata™ 20WG 0.6 oz



Check