



Tough Kyllinga Sedge Isn't So Tough

Celero® Herbicide is a selective, post-emergence herbicide for control of sedges including kyllinga in established residential and commercial turfgrass, golf courses and sod farms. Imazosulfuron, the active ingredient in Celero, is a systemic sulfonylurea (SU) herbicide that translocates down to the roots to deliver superior sedge control. Labeled for both broadcast and spot treatment in cool- and warm-season turf, Celero is the ideal solution for difficult-to-manage sedges.

COMPETITIVE ADVANTAGES

- Consistent post-emergence control of yellow, purple and annual sedges including green kyllinga
- Excellent cool- and warm-season turf tolerance
- Outstanding resistance management tool

AT-A-GLANCE INFO

EPA REG. NO.	59639-155
ACTIVE INGR.	imazosulfuron (75%)
FORMULATION	water dispersible granule
CHEM. FAMILY	sulfonylurea
HRAC NO.	2
SIGNAL WORD	caution
PACKAGE SIZE	4 x 1 lb



KEY USES

Established residential and commercial turfgrass (apartment complexes, golf courses, sod farms, roadsides, sports fields, campgrounds, office complexes, parks, parking areas, recreational sites, schools, residential lawns and commercial areas)

KEY WEEDS CONTROLLED

(See product label for complete list)

SEDGES

Nutsedge (purple, yellow)
Sedge (annual, cylindric, green kylinga)

BROADLEAF WEEDS

Burclover, California
Burweed, lawn
Chickweed (common, mouse-ear, sticky Gernaium, Carolina)
Henbit
Parsley-piert
Purslane, common

USE SUMMARY

Apply Celero® after nutsedge has reached the 3-leaf stage of growth. Only use a non-ionic surfactant that contains at least 80% active ingredient.

BROADCAST APPLICATION

RATE PER ACRE: 8-14 oz/20-30 gal + .25% v/v non-ionic surfactant (1 qt/100 gal). A second application may be applied 21 days after the initial treatment.

SPOT TREATMENT


RATE PER 1000 SQ FT: 0.25-0.33 oz/1-2 gal + 2 tsp non-ionic surfactant/gal of water. Occasionally shake the spray solution while spraying to ensure the spray solution remains well mixed. Spray the target weeds until the leaves are wet.

TANK MIXING

Celero® may be tank mixed with other herbicides registered for use in turfgrass for nutsedge and broad-spectrum broadleaf weed control.

CELERO DELIVERS EXCELLENT GREEN KYLLINGA CONTROL

Source: Fred Yelverton, North Carolina State University Applications made 6/3 and 6/24 to green kylinga.



PRECAUTIONS AND RESTRICTIONS

Do not apply through any type of irrigation system.

Do not apply by air.

Do not apply to golf course putting greens.

Do not irrigate within 4 hrs before or after application.

Do not apply if rain is expected within 4 hrs after application.

Do not mow turfgrass within 24 hrs after application.

Do not apply to moist or wet turfgrass (including dew).

Do not apply when daily high air temps are below 65°F or above 90°F.

Do not apply to turfgrass or nutsedge under stress due to drought, temperature, disease, low fertility, heavy thatch, mechanical injury or other stresses. Turf must be well established and receive at least two mowings before application.

Do not apply over-the-top to desirable flowers, ornamentals, vegetables, shrubs or trees.

Allow 4 weeks between application and seeding or sodding of turfgrass.

Do not apply more than 0.66 lb imazosulfuron/A (0.015 lb ai/1,000 sq ft) per application.



nufarm.com/usturf

©2026 Nufarm. For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usturf to download a full label. Important: Always read and follow label instructions. 16-TO-0019-D