

PrimeTime™

Weed Control Firepower for Every Season

PrimeTime™ Herbicide combines the firepower of four active ingredients to deliver outstanding and fast-acting post-emergence control of the toughest broadleaf weeds from early spring to late fall. Apply early season in spring on slow-growing weeds, use in summer for difficult-to-control weeds including goosegrass, and treat in fall for winter cleanup. Expect long-lasting control for cool- and warm-season lawns, athletic fields and golf courses.

COMPETITIVE ADVANTAGES

- Excellent weed control at suboptimal application timings provides one herbicide to use at any time of the growing season – spring, summer or fall
- Fast-acting broadleaf weed control with visible results within hours of application
- Consistent control of more than 250 labeled weeds
- Easy tank-mixing and handling
- Excellent turf tolerance for many warm- and cool-season turfgrasses
- Carfentrazone adds speed of visual injury and confidence of goosegrass and nimblewill control

PRIMETIME ACTIVES

2,4-D & DICAMBA + FLUROXYPYR +CARFENTRAZONE



Labeled for 250+ weeds



Difficult to Control Broadleaves



Annual Grasses

AT-A-GLANCE INFO

EPA REG. NO.	228-762
ACTIVE INGR.	2,4-D (1.6 lb/gal), dicamba (0.2 lb/gal), fluroxypyr (0.2 lb/gal), carfentrazone (0.04 lb/gal)
FORMULATION	liquid
CHEM. FAMILY	phenoxy, benzoic acid, pyridine, N-phenyl-triazolinones
HRAC NO.	4 and 14
SIGNAL WORD	warning
PACKAGE SIZE	12 x 1 qt tip-n-measure, 2 x 2.5 gal, 30 gal, 250 gal

nufarm

PrimeTime™

KEY USES

Lawns: commercial and residential
Parks and athletic fields
Golf courses*
Sod farms
Industrial areas: roadsides, rights-of-way, railroads, fence rows

*Not for use on bentgrass greens, tees and aprons

KEY WEEDS CONTROLLED

(See product label for complete list)

Bindweed
Bitterweed
Broomweed
Buckhorn
Burdock
Burweed
Buttercup
Carolina geranium
Carpetweed
Catsear
Chickweed
Clover
Cudweed
Dandelion
Dock
Dogbane
English daisy
Fiddleneck
Field pansy
Florida betony
Florida pusley
Goosegrass
Ground ivy
Hairy bittercress
Hawkweed
Henbit
Kochia
Knotweed
Lambsquarter
Mallow
Morning glory
Mouse-ear
Nimblewill
Oxalis
Parsley-piert
Pearlwort
Pennywort
Pigweed
Plantain
Poison ivy
Purslane
Rush
Sheep sorrel
Shepherds purse
Speedwell
Spotted spurge
Spurge
Thistle
Virginia buttonweed
Virginia pepperweed
Wild violet
Woodsorrel
Yarrow

USE SUMMARY

	RATE
COOL-SEASON TURF Incl. Bluegrass spp, Fescue spp, or Ryegrass spp. 4 pints/A is a good target for most cool season applications. Apply 5-6 pints/A for early spring applications when growth is slow, or for difficult-to-control weeds like wild violet or ground ivy.	4–6 pints/A
WARM-SEASON TURF* Bahiagrass, Bermudagrass, zoysiagrass *Do not use on St. Augustinegrass or carpetgrass.	2–4 pints/A
BENTGRASS* Creeping bentgrass Use reduced rates if grass is stressed from heat or drought. Difficult to control weeds may require repeat applications. On closely mowed turf, do not exceed 3 pints/A. Any slight yellowing of the creeping bentgrass turf will disappear after one week. Avoid swath overlaps to minimize grass injury and delay any second application for at least 4 weeks. *Do not use on bentgrass greens, tees or aprons.	2–4 pints/A
GOOSEGRASS Use the higher rate when goosegrass is mature. More mature goosegrass may require a repeat application for complete control – in these cases apply PrimeTime at 3-4 pts/A approximately 3 weeks apart. For Bermudagrass and zoysiagrass, apply a maximum of 3 pts/A. Slight yellowing may occur, but should disappear after 1 week.	3–5 pints/A

ELMORE, RUTGERS UNIVERSITY, 2022

PrimeTime was applied twice at 4 pts/A (July 15 and August 13) in a stand of perennial ryegrass where goosegrass had grown to a 1-tiller stage. Pylex® was applied once at 1.5 fl oz/A.

Both treatments provided 100% control of goosegrass.


Untreated


Pylex® Treated


PrimeTime Treated

nufarm

nufarm.com/usturf

©2026 Nufarm. For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usturf to download a full label. Important: Always read and follow label instructions. Pylex® is a registered trademark of BASF. 23-TO-0927-G