

# Product Information Sheet

Revised December. 2024



# DOCKMASTER

<b>Product</b>	<b>DOCKMASTER</b>			
<b>Active ingredient:</b>	660 g/l MCPA + 90 g/l dicamba as the dimethylamine salt			
<b>Formulation:</b>	Soluble concentrate (SL)			
<b>PCS number:</b>	06760			
<b>Mode of Action:</b>	Auxin mimic: HRAC group 4 (Legacy HRAC group O)			
<b>Pack size:</b>	5 litres			
<b>Aquatic buffer zone</b>	5 metres			
<b>Non-target Arthropod</b>	n/a			
<b>Benefits</b>	<ul style="list-style-type: none"><li>✓ Excellent control of docks &amp; chickweed – ideally suited to intensive dairy &amp; livestock sector.</li><li>✓ New-sown ley application.</li><li>✓ Short grazing and cutting interval – 7 days.</li><li>✓ Good fit for multi-cut silage &amp; rotational grazing.</li><li>✓ Convenient 5 litre pack size.</li></ul>			
<b>RECOMMENDATIONS</b>	Apply DOCKMASTER with a tractor mounted/trailed sprayer in good weather conditions with active plant growth			
<b>Crops</b>	<b>Full dose</b>	<b>Total dose</b>	<b>No. of application</b>	<b>Latest time of application</b>
New-sown ley	1.5 l/ha	1.5 l/ha	1 per 12 months	30 <sup>th</sup> September, minimum 7 days before cutting/mowing
Established grassland	1.5 l/ha	1.5 l/ha	1 per 12 months	30 <sup>th</sup> September, minimum 7 days before cutting/mowing

<b>Label weeds</b>	<b>Susceptible</b> <ul style="list-style-type: none"> <li>Broad leaf dock (<i>Rumex obtusifolius</i>)</li> <li>Common chickweed (<i>Stellaria media</i>)</li> <li>Corn buttercup (<i>Ranunculus arvensis</i>)</li> </ul>	<b>Moderately Susceptible</b> <ul style="list-style-type: none"> <li>Creeping thistle (<i>Cirsium arvense</i>)</li> <li>Creeping buttercup (<i>Ranunculus repens</i>)</li> <li>Dandelion (<i>Taraxacum officianale</i>)</li> <li>Common sorrel (<i>Rumex acetosa</i>)</li> </ul>
<b>Time of application:</b>	<b>Established Grassland:</b> Apply to actively growing weeds at leafy rosette stage. <b>New-sown ley:</b> Apply at minimum 4 leaf stage of grass and before 6 true leaves of weeds.	
<b>Wash-out procedure</b>	Wash equipment thoroughly immediately after use. Triple rinse sprayer, drain and allow to dry. Do not clean equipment near surface water and avoid contamination via drains from farmyards.	
<b>Agronomy Solutions</b>	<b>DOCKMASTER 1.5 l/ha</b> <i>A simple solution for excellent control of docks and chickweed in established grassland and new-sown ley</i>	
	<b>DOCKMASTER 1.5 l/ha + TOMAHAWK 0.75 l/ha</b> <i>A broad spectrum solution for new sown leys with excellent control of docks, chickweed, thistle, dandelion etc.</i>	
	<b>DOCKMASTER 1.5 l/ha + DEPITONE ULTRA 1.5 l/ha</b> <i>A useful mix on 6 month old or established grass for control of docks, buttercup, chickweed and thistle.</i>	
	<b>DOCKMASTER 1.5 l/ha + NU46 / AGRITOX 1.5 l/ha</b> <i>A useful mix for control of docks, thistle and buttercup in established grassland.</i>	
<b>Water Volume</b>	200-400 litres water/ha	
<b>Best Use Advice</b>	Apply DOCKMASTER in 200 litres water/ha to actively growing weeds. If ragwort is present in established grassland it is best to consider treatments based on incorporating full dose 2,4-D such as D50, Thrust or Lupo Ragwort is best treated in autumn for fields which are due to be cut in the following spring. Livestock should be excluded from treated areas until ragwort has died and become unpalatable.	
<b>Tank Mix</b>	Physically compatible with most herbicides, trace elements, insecticides & fungicides (check labels) – See nufarm.ie for latest tank mix list	
<b>Grazing Interval</b>	Do not graze for at least 7 days after spraying	
<b>Cutting Interval</b>	Do not mow or roll 4 days before or after application	
<b>Overseeding Interval</b>	No restriction on following crops	
<b>Clover Safety (Yes or No)</b>	No	
<b>Handheld Applicator (Yes or No)</b>	No	
<b>Following Crop Information</b>	No restriction on following crops	