

SENATOR® 600 FS

This seed has been treated with:

_____ mg imidacloprid/seed OR

_____ lbs imidacloprid/100 lbs of seed

CAUTION

TREATED SEED

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSE. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Treated seed may be harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Persons handling product, treating seed, or involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment must wear the following PPE: long-sleeved shirt, long pants, shoes and socks and chemical resistant gloves when handling treated seed.

ENVIRONMENTAL HAZARDS:

Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds.

USE RESTRICTIONS:

Treated seed must be planted into the soil at a depth greater than 1 inch.

The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lbs. Do not apply more than 0.5 lbs a.i. imidacloprid per acre per calendar year.

DO NOT graze or feed livestock on treated areas for 45 days after planting.

REPLANTING / CROP ROTATION RESTRICTIONS:

In the event of a crop failure or harvest of a crop grown from SENATOR 600 FS (imidacloprid) treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date SENATOR 600 FS (imidacloprid) treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid, tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

DISPOSAL:

Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate bodies of water when disposing of planting equipment wash water.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV020620

13718000

Senator is a Registered Trademark of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

